


Drilling Sequence Guidelines

Soft-Bone Protocol: Follow solid color bars on the surgical tray surface until the segmented color bar. The segmented color bar indicates the final drill for soft-bone protocol.

Dense-Bone Protocol: Follow solid color bars only. The last solid bar in the sequence represents the final drill for dense-bone.





Dríva[™] Gold Series Drills

0


21

NPMODG

snaps into TSVKITG

(TSV Surgical Kit)

- Recommended drill speed is 600-850 rpms
- design of the drill point. The 2.3 mmD Pilot Drill [TSV23G, TSV23SG] is the only drill that is close to the actual length (i.e., 8.0 mmL is actually 8.25 mmL).
- Surgical instruments are reusable up to 15 uses, excluding those labeled as single use. Surgical instruments are susceptible to damage and wear and should be inspected before use. The number of uses per drill will vary and depends on a variety of factors including bone density encountered, proper handling, and cleaning. Over time, repeat sterilizations may affect cutting efficiency and color appearance. Cutting edges should present a continuous edge and appear sharp. Check the latch-lock shank for wear to ensure the connection is not damaged. If inspection reveals signs of wear, damage, or unrecognizable color identification, replace the drill accordingly.



ZimVie DENTAL SOLUTIONS

The information provided in this guide serves as a quick reference only. Please refer to the Instructions for Use (IFU) for each product for complete information.

Implant Sizing and Placement Guidelines

Ensure the implant size and abutment angulation are appropriate for the occlusal load.

Eztetic Implants

- 3.1 mmD only, for anterior placement
- Should be splinted to additional implants when used in the pre-molar region and should not be used in the molar region

Tapered Screw-Vent[®] Implants

• 3.7, 4.1, 4.7, 6.0 mmD

Trabecular Metal Implants

- 3.7, 4.1, 4.7, 6.0 mmD
- 3.7 mmD Trabecular Metal Implants should be splinted to additional implants when used in the pre-molar region and should not be used in the molar region
- 4.1 mmD Trabecular Metal Implants should be splinted to additional implants when used in the molar region

Cleaning and Sterilization

Dríva Gold Series Drills and Kits with part numbers ending in the letter G utilize IFU "Cleaning and Sterilization of ZimVie Kits and Instruments" [P-ZBDINSTRP] available at ifu.biomet3i.com.

NOTE: These parts and kits have also been validated to the cleaning and sterilization guidelines in IFU 8874 "Instructions for Use for ZimVie Instrument Kit System and Dríva Drills" available at ifu.biomet3i.com.

Validated Cleaning/Disinfection Methods

Method	Description
Rigorous Manual Cleaning Instructions for Instruments	Enzymatic or alkaline soak and scrub followed by sonication
Rigorous Manual Cleaning Instructions for Trays	Enzymatic or alkaline soak and scrub
Combination Cleaning and Disinfection Instructions for Instruments	Enzymatic soak and scrub with sonication or alkaline soak with sonication followed by automated washer/disinfector cycle

Validated Sterilization Parameters

	Gravity Displacement Sterilizer (Full Cycle)	Pre-Vacuum Sterilizer (HI-VAC)
Catalog Number (Kit)	15 Minutes 132°C to 135°C (270°F to 275°F) 30 Minute Dry Time	4 Minute, 4 Pulse 132°C to 135°C (270°F to 275°F) 30 Minute Dry Time
TSVKITG, NPMODG	Х	x
Dríva Gold Series Drills and Stand-alone Instruments	Х	х

For more information, visit ZimVie.com

ZimVie 4555 Riverside Drive Palm Beach Gardens, FL 33410 1-800-342-5454 Phone: +1-561-776-6700 Fax: +1-561-776-1272


Unless otherwise indicated, as referenced herein, all trademarks and intellectual property rights are the property of ZimVie Inc. or an affiliate; and all products are manufactured by one or more of the dental subsidiaries of ZimVie Inc. (Biomet 3i, LLC, Zimmer Dental, Inc., etc.) and marketed and distributed by ZimVie Dental and its authorized marketing partners. For additional product information, please refer to the individual product labeling or instructions for use. Product clearance **CE** 2797 and availability may be limited to certain countries/regions. This material is intended for clinicians only and does not comprise medical advice or recommendations. Distribution to any other recipient is prohibited. This material may not be copied or reprinted without the express written consent of ZimVie. ZVINST0141 REV A 07/23 ©2023 ZimVie, Inc. All rights reserved.

Drilling Sequence

Eztetic Implants



Trabecular Metal and TSV Implants





ZimVie

Restoring Daily Life:













Bone Tap for placement in dense bone, 22 mml Drills shown, 16 mmL Drills have depth markings for placement of 8 and 10 mm implant lengths.





TRABECULAR METAL ONLY FOR DENSE BONE* TSV38G 4.1 mmD Cortical Bone Tap



OPTIONAL FOR DENSE BONE TT4 1 4.1 mmD Cortical Bone Tap



OPTIONAL FOR

DENSE BONE

4.7 mmD

Cortical Bone Tap

