

**Biologic Solutions** 

# InterGro® DBM Fibers

100% DBM cortical fibers ready for re-hydration with choice of fluid

# **InterGro DBM Fibers**

## **Bone Remodeling**

- 100% demineralized human cortical bone fibers offer osteoconductive and osteoinductive properties
- · Each lot is verified for osteoinductivity
- Fibers provide increased osteoinductivity when compared to the osteoinductivity of particulate formulations<sup>1</sup>



Re-hydrate with autologous blood products, saline, or antibiotic solution

Good Molding
Capability<sup>2</sup>
Co

### **Handling Characteristics**

- User-controlled graft consistency
- Flexible and moldable upon re-hydration

### Convenience

- All-inclusive bone grafting kit
- Available in 2cc, 5cc, or 10cc sizes



# Wicking Ability<sup>2\*</sup>



VS.





Dry

Re-hydrated

InterGro DBM Fibers

Cancellous Chips

## Instructions for Use\*\*

#### STEP 1

 Attach the 30cc vacuum syringe to the valve fitting on the side of the graft syringe containing DBM Fibers



- Pull the vacuum syringe plunger until fully extended
- Twist plunger to engage the locking mechanism
   Note: as the plunger is extended the air
   is being removed
- Twist off the 30cc vacuum syringe to remove

#### STEP 3

 Attach a dispensing syringe containing the liquid hydrating component of surgeon's choice onto the valve of the graft syringe



- o Ensure a minimum ratio of 0.6ml fluid to 1cc DBM fibers prior to attaching dispensing syringe to graft syringe
- Twist the dispensing syringe in a clockwise motion to allow fluid to be dispensed into the graft syringe

**Important Note:** The dispensing syringe is not included in the package of InterGro DBM Fibers. Utilize an off-the-shelf luer lock syringe to add fluid.



#### STEP 4

- · Detach the dispensing syringe and discard
- Pull back and forth for 10 seconds on the plunger of the graft syringe to fully immerse the graft material with fluid



#### STEP 5

- Remove the cap from the end of the graft syringe
- Push the plunger gently to remove graft material

<sup>\*</sup> Each 5cc sample was hydrated with 5 ml of PRP.

<sup>\*\*</sup> Please see package insert for complete instructions for use.

#### **Product Availability**

Product	Description	Product Number
00	InterGro DBM Fibers 2cc	CDF002
00	InterGro DBM Fibers 5cc	CDF005
0-8-1	InterGro DBM Fibers 10cc	CDF010

To order samples contact Zimmer Biomet Spine Marketing Department. InterGro DBM Fiber Marketing sample - CDF005S

#### References

- 1. Zimmer Biomet Internal Test Report TR-1020, Comparison of Osteoinductivity of DBM Powder to DBM Fiber
- 2. TR-1009, StaGraft Fiber DBM Bench Top Evaluation

#### **INDICATIONS**

InterGro DBM Fibers can be used to fill bony voids or gaps that have been surgically created, or for filling osseous defects in nonweight bearing applications.

InterGro DBM Fibers may be used with orthopedic, spinal, reconstructive, craniofacial, maxillofacial and periodontal bone grafting procedures. It can be used alone or in combination with autologous bone, or other forms of allogeneic bone in grafting procedures of non-weight bearing value. It can be used or hydrated with autologous blood, bone marrow aspirate, or autologous blood derived products such as platelet rich plasma and platelet poor plasma. It may also be hydrated with saline or antibiotic solution.

All content herein is protected by copyright, trademarks and other intellectual property rights owned by or licensed to Zimmer Biomet or its affiliates unless otherwise indicated, and must not be redistributed, duplicated or disclosed, in whole or in part, without the express written consent of Zimmer Biomet.

This material is intended for health care professionals and the Zimmer Biomet sales force. Distribution to any other recipient is prohibited. For complete product information, including indications, contraindications, warnings, precautions, and potential adverse effects, see the package insert and www.zimmerbiomet.com.

©2018 Zimmer Biomet Spine



